

Rank Polynomials of Fence Posets are Unimodal

(joint work with Mohan Ravichandran)

Ezgi KANTARCI OĞUZ

Boğaziçi University
İstanbul, Turkey


March 23, 2022

What are fences?

Let $\alpha = (\alpha_1, \alpha_2, \dots, \alpha_s)$ be a composition of n . The fence poset of α , denoted $F(\alpha)$ is the poset on x_1, x_2, \dots, x_{n+1} with the order relations:

$$x_1 \preceq x_2 \preceq \dots \preceq x_{\alpha_1+1} \succeq x_{\alpha_1+2} \succeq \dots \succeq x_{\alpha_1+\alpha_2+1} \preceq x_{\alpha_1+\alpha_2+2} \preceq \dots$$

Example ($\alpha = (2, 1, 1, 3)$)



For a composition of n , we get a poset of $n + 1$ nodes.


An **ideal** of a fence is a down-closed subset: $x \in I, y \preceq x \Rightarrow y \in I$.

$$\#I = \text{rank}(I)$$

An **ideal** of a fence is a down-closed subset: $x \in I, y \preceq x \Rightarrow y \in I$.

$$\#I = \text{rank}(I)$$


Example ($\alpha = (2, 1, 1, 3)$)



An **ideal** of a fence is a down-closed subset: $x \in I, y \preceq x \Rightarrow y \in I$.

$$\#I = \text{rank}(I)$$

Example ($\alpha = (2, 1, 1, 3)$)




1 ideal of rank 0,

An **ideal** of a fence is a down-closed subset: $x \in I, y \preceq x \Rightarrow y \in I$.

$$\#I = \text{rank}(I)$$

Example ($\alpha = (2, 1, 1, 3)$)




1 ideal of rank 0,

An **ideal** of a fence is a down-closed subset: $x \in I, y \preceq x \Rightarrow y \in I$.

$$\#I = \text{rank}(I)$$

Example ($\alpha = (2, 1, 1, 3)$)




1 ideal of rank 0,

An **ideal** of a fence is a down-closed subset: $x \in I, y \preceq x \Rightarrow y \in I$.

$$\#I = \text{rank}(I)$$

Example $(\alpha = (2, 1, 1, 3))$




1 ideal of rank 0, 3 ideals of rank 1,

An **ideal** of a fence is a down-closed subset: $x \in I, y \preceq x \Rightarrow y \in I$.

$$\#I = \text{rank}(I)$$

Example ($\alpha = (2, 1, 1, 3)$)




1 ideal of rank 0, 3 ideals of rank 1,

An **ideal** of a fence is a down-closed subset: $x \in I, y \preceq x \Rightarrow y \in I$.

$$\#I = \text{rank}(I)$$

Example ($\alpha = (2, 1, 1, 3)$)




1 ideal of rank 0, 3 ideals of rank 1,

An **ideal** of a fence is a down-closed subset: $x \in I, y \preceq x \Rightarrow y \in I$.

$$\#I = \text{rank}(I)$$

Example ($\alpha = (2, 1, 1, 3)$)




1 ideal of rank 0, 3 ideals of rank 1,

An **ideal** of a fence is a down-closed subset: $x \in I, y \preceq x \Rightarrow y \in I$.

$$\#I = \text{rank}(I)$$

Example ($\alpha = (2, 1, 1, 3)$)




1 ideal of rank 0, 3 ideals of rank 1,

An **ideal** of a fence is a down-closed subset: $x \in I, y \preceq x \Rightarrow y \in I$.

$$\#I = \text{rank}(I)$$

Example $(\alpha = (2, 1, 1, 3))$




1 ideal of rank 0, 3 ideals of rank 1, 5 ideals of rank 2, ...

An **ideal** of a fence is a down-closed subset: $x \in I, y \preceq x \Rightarrow y \in I$.

$$\#I = \text{rank}(I)$$

Example ($\alpha = (2, 1, 1, 3)$)




1 ideal of rank 0, 3 ideals of rank 1, 5 ideals of rank 2, ...

$(1, 3, 5, 6, 6, 5, 3, 2, 1) \leftarrow$ Rank sequence.

An **ideal** of a fence is a down-closed subset: $x \in I, y \preceq x \Rightarrow y \in I$.

$$\#I = \text{rank}(I)$$

Example ($\alpha = (2, 1, 1, 3)$)



1 ideal of rank 0, 3 ideals of rank 1, 5 ideals of rank 2, ...

$(1, 3, 5, 6, 6, 5, 3, 2, 1) \leftarrow$ Rank sequence.

$1 + 3q + 5q^2 + 6q^3 + 6q^4 + 5q^5 + 3q^6 + 2q^7 + q^8 \leftarrow$ Rank polynomial.

A q -deformation for rational numbers

Recently, a q -deformation rational numbers was introduced by Morier-Genoud and Ovsienko¹. Their definition has a *convergence* property, which allows us to extend them to real numbers.

¹Morier-Genoud and Ovsienko, “ q -deformed rationals and q -continued fractions”.

A q -deformation for rational numbers

Recently, a q -deformation rational numbers was introduced by Morier-Genoud and Ovsienko¹. Their definition has a *convergence* property, which allows us to extend them to real numbers.

For a given rational number r/s , we first write it as a continued fraction.

$$\frac{r}{s} = a_1 + \frac{1}{a_2 + \frac{1}{a_3 + \frac{1}{\ddots + \frac{1}{a_{2m}}}}} = c_1 - \frac{1}{c_2 - \frac{1}{c_3 - \frac{1}{\ddots - \frac{1}{c_k}}}}$$

$$a_i \in \mathbb{Z}, a_i \geq 1 \text{ for } i \geq 2$$

$$c_i \in \mathbb{Z}, c_i \geq 2 \text{ for } i \geq 2$$

¹Morier-Genoud and Ovsienko, “ q -deformed rationals and q -continued fractions”.

A q -deformation for rational numbers

Then we replace the expansion terms with q -integers (q^{-1} -integers for a_{2k}), and the 1's with powers of q .

$$\left[\frac{r}{s} \right]_q := [a_1]_q + \frac{q^{a_1}}{[a_2]_{q^{-1}} + \frac{q^{-a_2}}{\dots + \frac{q^{a_{2m-1}}}{[a_{2m}]_{q^{-1}}}}} = [c_1]_q - \frac{q^{c_1-1}}{[c_2]_q - \frac{q^{c_2-1}}{\dots - \frac{q^{c_{k-1}-1}}{[c_k]_q}}}$$

A q -deformation for rational numbers

Then we replace the expansion terms with q -integers (q^{-1} -integers for a_{2k}), and the 1's with powers of q .

$$\left[\frac{r}{s} \right]_q := [a_1]_q + \frac{q^{a_1}}{[a_2]_{q^{-1}} + \frac{q^{-a_2}}{\dots + \frac{q^{a_{2m-1}}}{[a_{2m}]_{q^{-1}}}}} = [c_1]_q - \frac{q^{c_1-1}}{[c_2]_q - \frac{q^{c_2-1}}{\dots - \frac{q^{c_{k-1}-1}}{[c_k]_q}}}$$

A cool thing: The two expressions give the same q -deformation.

A q -deformation for rational numbers

Then we replace the expansion terms with q -integers (q^{-1} -integers for a_{2k}), and the 1's with powers of q .

$$\left[\frac{r}{s} \right]_q := [a_1]_q + \frac{q^{a_1}}{[a_2]_{q^{-1}} + \frac{q^{-a_2}}{\dots + \frac{q^{a_{2m-1}}}{[a_{2m}]_{q^{-1}}}} = [c_1]_q - \frac{q^{c_1-1}}{[c_2]_q - \frac{q^{c_2-1}}{\dots - \frac{q^{c_{k-1}-1}}{[c_k]_q}}$$

A cool thing: The two expressions give the same q -deformation.

Another cool thing: $\left[\frac{r}{s} \right]_q = \frac{R(q)}{S(q)}$ where $R(q), S(q) \in \mathbb{Z}[q]$ are polynomials that evaluate to r and s respectively.

A q -deformation for rational numbers

Then we replace the expansion terms with q -integers (q^{-1} -integers for a_{2k}), and the 1's with powers of q .

$$\left[\frac{r}{s} \right]_q := [a_1]_q + \frac{q^{a_1}}{[a_2]_{q^{-1}} + \frac{q^{-a_2}}{\dots + \frac{q^{a_{2m-1}}}{[a_{2m}]_{q^{-1}}}} = [c_1]_q - \frac{q^{c_1-1}}{[c_2]_q - \frac{q^{c_2-1}}{\dots - \frac{q^{c_{k-1}-1}}{[c_k]_q}}$$

A cool thing: The two expressions give the same q -deformation.

Another cool thing: $\left[\frac{r}{s} \right]_q = \frac{R(q)}{S(q)}$ where $R(q), S(q) \in \mathbb{Z}[q]$ are polynomials that evaluate to r and s respectively.

Also, when $\frac{r}{s} \geq 0$ the coefficients are non-negative.

Example

$$\frac{32}{9} = 3 + \frac{1}{1 + \frac{1}{1 + \frac{1}{4}}} = 4 - \frac{1}{3 - \frac{1}{2 - \frac{1}{2 - \frac{1}{2}}}}$$

Example

$$\frac{32}{9} = 3 + \frac{1}{1 + \frac{1}{1 + \frac{1}{4}}} = 4 - \frac{1}{3 - \frac{1}{2 - \frac{1}{2 - \frac{1}{2}}}}$$

$$\left[\frac{32}{9} \right]_q = [3]_q + \frac{q^3}{[1]_{q^{-1}} + \frac{q}{[4]_{q^{-1}}}} = [4]_q - \frac{q^4}{[3]_q - \frac{q^3}{[2]_q - \frac{q^2}{[2]_q - \frac{q^2}{[2]_q}}]}$$

Example

$$\frac{32}{9} = 3 + \frac{1}{1 + \frac{1}{1 + \frac{1}{4}}} = 4 - \frac{1}{3 - \frac{1}{2 - \frac{1}{2 - \frac{1}{2}}}}$$

$$\left[\frac{32}{9} \right]_q = [3]_q + \frac{q^3}{[1]_{q^{-1}} + \frac{q^{-1}}{[1]_q + \frac{q}{[4]_{q^{-1}}}}} = [4]_q - \frac{q^4}{[3]_q - \frac{q^3}{[2]_q - \frac{q^2}{[2]_q - \frac{q^2}{[2]_q}}]}$$

$$\left[\frac{32}{9} \right]_q = \frac{1 + 3q + 5q^2 + 6q^3 + 6q^4 + 5q^5 + 3q^6 + 2q^7 + q^8}{1 + 2q + 2q^2 + 2q^3 + q^4 + q^5}.$$

Example

$$\left[\frac{32}{9} \right]_q = [3]_q + \frac{q^3}{[1]_{q^{-1}} + \frac{q^{-1}}{[1]_q + \frac{q}{[4]_{q^{-1}}}}} = [4]_q - \frac{q^4}{[3]_q - \frac{q^3}{[2]_q - \frac{q^2}{[2]_q - \frac{q^2}{[2]_q}}]}$$

$$\left[\frac{r}{s} \right]_q = \frac{\text{Rank polynomial for } (2, 1, 1, 3)}{\text{Rank polynomial for } (1, 3)}$$

Example

$$\left[\frac{32}{9} \right]_q = [3]_q + \frac{q^3}{[1]_{q^{-1}} + \frac{q^{-1}}{[1]_q + \frac{q}{[4]_{q^{-1}}}}} = [4]_q - \frac{q^4}{[3]_q - \frac{q^3}{[2]_q - \frac{q^2}{[2]_q - \frac{q^2}{[2]_q}}]}$$

$$\left[\frac{r}{s} \right]_q = \frac{\text{Rank polynomial for } (2, 1, 1, 3)}{\text{Rank polynomial for } (1, 3)}$$

In general, if r/s corresponds to $[a_1, a_2, \dots, a_{2m}]$, we have

$$\left[\frac{r}{s} \right]_q = \frac{\text{Rank polynomial for } (a_1 - 1, a_2, a_3, \dots, a_{2m} - 1)}{\text{Rank polynomial for } (0, a_2 - 1, a_3, \dots, a_{2m} - 1)}$$

A closer look at rank sequences for fences

$$\begin{aligned}(2, 1, 1, 3) &\rightarrow (1, 3, 5, 6, 6, 5, 3, 2, 1) \\(3, 1, 1, 2) &\rightarrow (1, 2, 3, 5, 6, 6, 5, 3, 1) \\(1, 2, 1, 3) &\rightarrow (1, 3, 5, 6, 6, 5, 4, 2, 1) \\(1, 1, 2, 3) &\rightarrow (1, 3, 5, 7, 7, 5, 4, 2, 1) \\(2, 2, 3) &\rightarrow (1, 2, 4, 5, 6, 6, 4, 2, 1) \\(2, 3, 2) &\rightarrow (1, 2, 4, 6, 7, 6, 4, 2, 1) \\(2, 1, 4) &\rightarrow (1, 2, 3, 3, 4, 4, 3, 2, 1) \\(2, 1, 2, 1, 1) &\rightarrow (1, 3, 6, 7, 8, 7, 5, 3, 1)\end{aligned}$$

A closer look at rank sequences for fences

$$\begin{aligned}(2, 1, 1, 3) &\rightarrow (1, 3, 5, 6, 6, 5, 3, 2, 1) \\(3, 1, 1, 2) &\rightarrow (1, 2, 3, 5, 6, 6, 5, 3, 1) \\(1, 2, 1, 3) &\rightarrow (1, 3, 5, 6, 6, 5, 4, 2, 1) \\(1, 1, 2, 3) &\rightarrow (1, 3, 5, 7, 7, 5, 4, 2, 1) \\(2, 2, 3) &\rightarrow (1, 2, 4, 5, 6, 6, 4, 2, 1) \\(2, 3, 2) &\rightarrow (1, 2, 4, 6, 7, 6, 4, 2, 1) \\(2, 1, 4) &\rightarrow (1, 2, 3, 3, 4, 4, 3, 2, 1) \\(2, 1, 2, 1, 1) &\rightarrow (1, 3, 6, 7, 8, 7, 5, 3, 1)\end{aligned}$$

Conjecture (Morier-Genoud, Ovsienko, 2020)

The rank polynomials of fence posets are unimodal.

What more can we say?

Consider $(2, 1, 1, 3) \rightarrow (1, 3, 5, 6, 6, 5, 3, 2, 1)$.

What more can we say?

Consider $(2, 1, 1, 3) \rightarrow (1, 3, 5, 6, 6, 5, 3, 2, 1)$.

We have $1 \leq 1 \leq 2 \leq 3 \leq 3 \leq 5 \leq 5 \leq 6 \leq 6$.

We call such a sequence **bottom-interlacing**:

$$a_n \leq a_0 \leq a_{n-1} \leq a_1 \leq \dots \leq a_{\lfloor n/2 \rfloor}. \quad (\text{BI})$$

What more can we say?

Consider $(2, 1, 1, 3) \rightarrow (1, 3, 5, 6, 6, 5, 3, 2, 1)$.

We have $1 \leq 1 \leq 2 \leq 3 \leq 3 \leq 5 \leq 5 \leq 6 \leq 6$.

We call such a sequence **bottom-interlacing**:

$$a_n \leq a_0 \leq a_{n-1} \leq a_1 \leq \dots \leq a_{\lfloor n/2 \rfloor}. \quad (\text{BI})$$

We call similarly have **top-interlacing** sequences:

$$a_0 \leq a_n \leq a_1 \leq a_{n-1} \leq \dots \leq a_{\lceil n/2 \rceil}. \quad (\text{TI})$$

What more can we say?

Consider $(2, 1, 1, 3) \rightarrow (1, 3, 5, 6, 6, 5, 3, 2, 1)$.

We have $1 \leq 1 \leq 2 \leq 3 \leq 3 \leq 5 \leq 5 \leq 6 \leq 6$.

We call such a sequence **bottom-interlacing**:

$$a_n \leq a_0 \leq a_{n-1} \leq a_1 \leq \dots \leq a_{\lfloor n/2 \rfloor}. \quad (\text{BI})$$

We call similarly have **top-interlacing** sequences:

$$a_0 \leq a_n \leq a_1 \leq a_{n-1} \leq \dots \leq a_{\lfloor n/2 \rfloor}. \quad (\text{TI})$$

For example, the rank sequence $(1, 2, 4, 5, 6, 6, 4, 2, 1)$ of $(2, 2, 3)$ is top interlacing:

$$1 \leq 1 \leq 2 \leq 2 \leq 4 \leq 4 \leq 5 \leq 6 \leq 6.$$

What more can we say?

- $(2, 1, 1, 3) \rightarrow (1, 3, 5, 6, 6, 5, 3, 2, 1) \rightarrow \text{BI}$
- $(3, 1, 1, 2) \rightarrow (1, 3, 5, 6, 6, 5, 3, 2, 1) \rightarrow \text{BI}$
- $(1, 2, 1, 3) \rightarrow (1, 3, 5, 6, 6, 5, 4, 2, 1) \rightarrow \text{BI}$
- $(1, 1, 2, 3) \rightarrow (1, 3, 5, 7, 7, 5, 4, 2, 1) \rightarrow \text{BI}$
- $(2, 2, 3) \rightarrow (1, 2, 4, 5, 6, 6, 4, 2, 1) \rightarrow \text{TI}$
- $(2, 3, 2) \rightarrow (1, 2, 4, 6, 7, 6, 4, 2, 1) \rightarrow \text{BI, TI (symmetric)}$
- $(2, 1, 4) \rightarrow (1, 2, 3, 3, 4, 4, 3, 2, 1) \rightarrow \text{TI}$
- $(2, 1, 2, 1, 1) \rightarrow (1, 3, 6, 7, 8, 7, 5, 3, 1) \rightarrow \text{BI}$

What more can we say?

$$(2, 1, 1, 3) \rightarrow (1, 3, 5, 6, 6, 5, 3, 2, 1) \rightarrow \text{BI}$$

$$(3, 1, 1, 2) \rightarrow (1, 3, 5, 6, 6, 5, 3, 2, 1) \rightarrow \text{BI}$$

$$(1, 2, 1, 3) \rightarrow (1, 3, 5, 6, 6, 5, 4, 2, 1) \rightarrow \text{BI}$$

$$(1, 1, 2, 3) \rightarrow (1, 3, 5, 7, 7, 5, 4, 2, 1) \rightarrow \text{BI}$$

$$(2, 2, 3) \rightarrow (1, 2, 4, 5, 6, 6, 4, 2, 1) \rightarrow \text{TI}$$

$$(2, 3, 2) \rightarrow (1, 2, 4, 6, 7, 6, 4, 2, 1) \rightarrow \text{BI, TI (symmetric)}$$

$$(2, 1, 4) \rightarrow (1, 2, 3, 3, 4, 4, 3, 2, 1) \rightarrow \text{TI}$$

$$(2, 1, 2, 1, 1) \rightarrow (1, 3, 6, 7, 8, 7, 5, 3, 1) \rightarrow \text{BI}$$

Conjecture (McConville, Sagan, Smyth, 2021²)


Suppose $\alpha = (\alpha_1, \alpha_2, \dots, \alpha_s)$.

- (a) If $s = 1$ then $r(\alpha) = (1, 1, \dots, 1)$ is symmetric.
- (b) If s is even, then $r(\alpha)$ is bottom interlacing.
- (c) If $s \geq 3$ is odd we have:
 - (i) If $\alpha_1 > \alpha_s$ then $r(\alpha)$ is bottom interlacing.
 - (ii) If $\alpha_1 < \alpha_s$ then $r(\alpha)$ is top interlacing.
 - (iii) If $\alpha_1 = \alpha_s$ then $r(\alpha)$ is symmetric, bottom interlacing, or top interlacing depending on whether $r(\alpha_2, \alpha_3, \dots, \alpha_{s-1})$ is symmetric, top interlacing, or bottom interlacing, respectively.

²McConville, B. E. Sagan, and Smyth, *On a rank-unimodality conjecture of Morier-Genoud and Ovsienko*.


What if we close up the fence?

Example ($\alpha = (2, 1, 1, 3)$)




What if we close up the fence?

Example ($\alpha = (2, 1, 1, 3)$)



What if we close up the fence?


Example ($\alpha = (2, 1, 1, 3)$)



The *circular* fence has rank sequence $(1, 2, 3, 4, 4, 3, 2, 1)$.

What if we close up the fence?

Example ($\alpha = (2, 1, 1, 3)$)




The *circular* fence has rank sequence $(1, 2, 3, 4, 4, 3, 2, 1)$.

It is symmetric. Is this always so?

What if we close up the fence?

Example ($\alpha = (2, 1, 1, 3)$)



The *circular* fence has rank sequence $(1, 2, 3, 4, 4, 3, 2, 1)$.

It is symmetric. Is this always so?

Answer: Yes, but it is not trivial to prove.

Theorem (Kantarci Oğuz, Ravichandran, 2021³)

Rank polynomials of circular fence posets are symmetric.


³Kantarci Oğuz and Ravichandran, *Rank Polynomials of Fence Posets are Unimodal.*

⁴Elizalde and B. Sagan, *Partial rank symmetry of distributive lattices for fences.*

Rank polynomials of circular fence posets are symmetric.

Our proof:

We have one case that is trivially symmetric: $(k, 1, 1, \dots, 1)$.



We show that moving a node from one segment to the next does not break symmetry.


³Kantarci Oğuz and Ravichandran, *Rank Polynomials of Fence Posets are Unimodal*.

⁴Elizalde and B. Sagan, *Partial rank symmetry of distributive lattices for fences*.

Rank polynomials of circular fence posets are symmetric.

Our proof:

We have one case that is trivially symmetric: $(k, 1, 1, \dots, 1)$.



We show that moving a node from one segment to the next does not break symmetry.

>> Recent bijective proof by Sagan and Elizalde⁴.

³Kantarıcı Oğuz and Ravichandran, *Rank Polynomials of Fence Posets are Unimodal*.

⁴Elizalde and B. Sagan, *Partial rank symmetry of distributive lattices for fences*.



The next step

There are several natural ways to associate a circular fence to a given fence.

The next step

There are several natural ways to associate a circular fence to a given fence.



Example (Adding the relation $x_1 \succeq x_8$)



The next step

There are several natural ways to associate a circular fence to a given fence.

Example (Adding the relation $x_1 \succeq x_8$)





$$\sum_I q^{\text{rank}(I)} = \sum_{\{I \mid x_1 \in I \Rightarrow x_8 \in I\}} q^{\text{rank}(I)} + \sum_{\{I \mid x_1 \in I, x_8 \notin I\}} q^{\text{rank}(I)}$$

The next step

There are several natural ways to associate a circular fence to a given fence.

Example (Adding the relation $x_1 \succeq x_8$)





$$\sum_I q^{\text{rank}(I)} = \sum_{\{I \mid x_1 \in I \Rightarrow x_8 \in I\}} q^{\text{rank}(I)} + \sum_{\{I \mid x_1 \in I, x_8 \notin I\}} q^{\text{rank}(I)}$$

The next step

There are several natural ways to associate a circular fence to a given fence.

Example (Adding the relation $x_1 \succeq x_8$)





$$\sum_I q^{\text{rank}(I)} = \sum_{\{I \mid x_1 \in I \Rightarrow x_8 \in I\}} q^{\text{rank}(I)} + \sum_{\{I \mid x_1 \in I, x_8 \notin I\}} q^{\text{rank}(I)}$$

The next step

There are several natural ways to associate a circular fence to a given fence.

Example (Adding the relation $x_1 \succeq x_8$)



$$\sum_I q^{\text{rank}(I)} = \sum_{\{I \mid x_1 \in I \Rightarrow x_8 \in I\}} q^{\text{rank}(I)} + \sum_{\{I \mid x_1 \in I, x_8 \notin I\}} q^{\text{rank}(I)}$$



circular rank
polynomial

$q \times$ rank polynomial
for $(1, 1)$

The next step

There are several natural ways to associate a circular fence to a given fence.

Example (Adding the relation $x_1 \succeq x_8$)



$$\sum_I q^{\text{rank}(I)} = \sum_{\{I \mid x_1 \in I \Rightarrow x_8 \in I\}} q^{\text{rank}(I)} + \sum_{\{I \mid x_1 \in I, x_8 \notin I\}} q^{\text{rank}(I)}$$

circular rank
polynomial
(*symmetric*)

$q \times$ rank polynomial
for $(1, 1)$
(*smaller, shifted center*)

What does this tell us about the rank polynomial?

$$\begin{array}{lll} \text{symmetric piece} & (1, 2, 3, 5, 5, 5, 3, 2, 1) & b_0 = b_n, b_1 = b_{n-1}, \dots \\ + & + & \\ \text{smaller piece,} & (0, 1, 2, 1, 1, 0, 0, 0, 0) & c_0 \geq c_n, c_1 \geq c_{n-1}, \dots \\ \text{shifted center} & & \\ = & = & \\ \sum_I q^{\text{rank}(I)} & (1, 3, 5, 6, 6, 5, 3, 2, 1) & a_0 \geq a_n, a_1 \geq a_{n-1}, \dots \end{array}$$

What does this tell us about the rank polynomial?

$$\begin{array}{lll} \text{symmetric piece} & (1, 2, 3, 5, 5, 5, 3, 2, 1) & b_0 = b_n, b_1 = b_{n-1}, \dots \\ + & + & \\ \text{smaller piece,} & (0, 1, 2, 1, 1, 0, 0, 0, 0) & c_0 \geq c_n, c_1 \geq c_{n-1}, \dots \\ \text{shifted center} & & \\ = & = & \\ \sum_I q^{\text{rank}(I)} & (1, 3, 5, 6, 6, 5, 3, 2, 1) & a_0 \geq a_n, a_1 \geq a_{n-1}, \dots \end{array}$$

This gives us half of the equations for being bottom interlacing:

$$a_n \leq a_0, \quad a_{n-1} \leq a_1, \quad a_{n-2} \leq a_2, \quad a_{n-3} \leq a_3, \dots$$

What does this tell us about the rank polynomial?

$$\begin{array}{lll} \text{symmetric piece} & (1, 2, 3, 5, 5, 5, 3, 2, 1) & b_0 = b_n, b_1 = b_{n-1}, \dots \\ + & + & \\ \text{smaller piece,} & (0, 1, 2, 1, 1, 0, 0, 0, 0) & c_0 \geq c_n, c_1 \geq c_{n-1}, \dots \\ \text{shifted center} & & \\ = & = & \\ \sum_I q^{\text{rank}(I)} & (1, 3, 5, 6, 6, 5, 3, 2, 1) & a_0 \geq a_n, a_1 \geq a_{n-1}, \dots \end{array}$$

This gives us half of the equations for being bottom interlacing:

$$a_n \leq a_0, \quad a_{n-1} \leq a_1, \quad a_{n-2} \leq a_2, \quad a_{n-3} \leq a_3, \dots$$

$$a_n \leq a_0 \leq a_{n-1} \leq a_1 \leq a_{n-2} \leq a_2 \leq a_{n-3} \leq a_3 \leq \dots \quad (\text{BI})$$

What does this tell us about the rank polynomial?

$$\begin{array}{lll} \text{symmetric piece} & (1, 2, 3, 5, 5, 5, 3, 2, 1) & b_0 = b_n, b_1 = b_{n-1}, \dots \\ + & + & \\ \text{smaller piece,} & (0, 1, 2, 1, 1, 0, 0, 0, 0) & c_0 \geq c_n, c_1 \geq c_{n-1}, \dots \\ \text{shifted center} & & \\ = & = & \\ \sum_I q^{\text{rank}(I)} & (1, 3, 5, 6, 6, 5, 3, 2, 1) & a_0 \geq a_n, a_1 \geq a_{n-1}, \dots \end{array}$$

This gives us half of the equations for being bottom interlacing:

$$a_n \leq a_0, \quad a_{n-1} \leq a_1, \quad a_{n-2} \leq a_2, \quad a_{n-3} \leq a_3, \dots$$


$$a_n \leq a_0 \leq a_{n-1} \leq a_1 \leq a_{n-2} \leq a_2 \leq a_{n-3} \leq a_3 \leq \dots \quad (\text{BI})$$

We need a way to shift the pairings to $(a_0, a_{n-1}), (a_1, a_{n+1}), \dots$ to get the rest of the inequalities.

Let us associate another circular fence to our fence.


Let us associate another circular fence to our fence.

Example (Connecting x_8 and x_1 by a minimal node x_0)



Let us associate another circular fence to our fence.


Example (Connecting x_8 and x_1 by a minimal node x_0)



$$\sum_{\{I \mid x_0 \in I\}} q^{\text{rank}(I)} = \sum_I q^{\text{rank}(I)} - \sum_{\{I \mid x_0 \notin I\}} q^{\text{rank}(I)}$$

Let us associate another circular fence to our fence.

Example (Connecting x_8 and x_1 by a minimal node x_0)



$$\sum_{\{I|x_0 \in I\}} q^{\text{rank}(I)} = \sum_I q^{\text{rank}(I)} - \sum_{\{I|x_0 \notin I\}} q^{\text{rank}(I)}$$


$q \times \text{rank}$
polynomial for $(2, 1, 1, 3)$

circular rank
polynomial

rank polynomial
for (0)

Let us associate another circular fence to our fence.

Example (Connecting x_8 and x_1 by a minimal node x_0)



$$\sum_{\{I | x_0 \in I\}} q^{\text{rank}(I)} = \sum_I q^{\text{rank}(I)} - \sum_{\{I | x_0 \notin I\}} q^{\text{rank}(I)}$$

$q \times \text{rank}$
polynomial for $(2, 1, 1, 3)$

circular rank
polynomial
(*symmetric,*
shifted center)

rank polynomial
for (0)
(*smaller,*
shifted center)

On the rank polynomial side

symmetric piece larger	(1, 2, 3, 5, 6, 6, 5, 3, 2, 1)	$b_0 = b_{n+1}, b_1 = b_n, \dots$
—	—	
smaller piece, shifted center	(1, 1, 0, 0, 0, 0, 0, 0, 0)	$c_0 \geq c_n, c_1 \geq c_{n-1}, \dots$
=	=	
(0, a_0, a_1, \dots, a_n)	(0, 1, 3, 5, 6, 6, 5, 3, 2, 1)	$0 \leq a_n, a_0 \leq a_{n-1} \dots$

On the rank polynomial side

$$\begin{array}{rcc}
 \text{symmetric piece} & (1, 2, 3, 5, 6, 6, 5, 3, 2, 1) & b_0 = b_{n+1}, b_1 = b_n, \dots \\
 \text{larger} & & \\
 - & - & \\
 \text{smaller piece,} & (1, 1, 0, 0, 0, 0, 0, 0, 0) & c_0 \geq c_n, c_1 \geq c_{n-1}, \dots \\
 \text{shifted center} & & \\
 = & = & \\
 (0, a_0, a_1, \dots, a_n) & (0, 1, 3, 5, 6, 6, 5, 3, 2, 1) & 0 \leq a_n, a_0 \leq a_{n-1} \dots
 \end{array}$$

This gives us the other half of the bottom-interlacing equations:

$$\begin{array}{c}
 a_n \leq a_0, \quad a_{n-1} \leq a_1, \quad a_{n-2} \leq a_2, \quad a_{n-3} \leq a_3, \dots \\
 + \\
 a_0 \leq a_{n-1}, \quad a_1 \leq a_{n-2}, \quad a_2 \leq a_{n-3}, \dots \\
 = \\
 a_n \leq a_0 \leq a_{n-1} \leq a_1 \leq a_{n-2} \leq a_2 \leq a_{n-3} \leq a_3 \leq \dots \quad (\text{BI})
 \end{array}$$

Theorem (Kantarci Oğuz, Ravichandran, 2021)

Rank polynomials of fence posets are unimodal.

In particular, for $\alpha = (\alpha_1, \alpha_2, \dots, \alpha_s)$ we have:

- (a) If $s = 1$ then $r(\alpha) = (1, 1, \dots, 1)$ is symmetric.*
- (b) If s is even, then $r(\alpha)$ is bottom interlacing.*
- (c) If $s \geq 3$ is odd we have:
 - (i) If $\alpha_1 > \alpha_s$ then $r(\alpha)$ is bottom interlacing.*
 - (ii) If $\alpha_1 < \alpha_s$ then $r(\alpha)$ is top interlacing.*
 - (iii) If $\alpha_1 = \alpha_s$ then $r(\alpha)$ is symmetric, bottom interlacing, or top interlacing depending on whether $r(\alpha_2, \alpha_3, \dots, \alpha_{s-1})$ is symmetric, top interlacing, or bottom interlacing, respectively.**

What about the rank polynomials of circular fence posets?

Are they also unimodal?

What about the rank polynomials of circular fence posets?

Are they also unimodal? **Answer:** Not always.

For the circular poset $(1, a, 1, a)$ we get a small dip in the middle:

$$(1, 2, \dots, a, a + 1, a, a + 1, a, a - 1, \dots, 2, 1).$$

What about the rank polynomials of circular fence posets?

Are they also unimodal? **Answer:** Not always.

For the circular poset $(1, a, 1, a)$ we get a small dip in the middle:

$$(1, 2, \dots, a, a + 1, a, a + 1, a, a - 1, \dots, 2, 1).$$

Nicer answer: Almost always.

What about the rank polynomials of circular fence posets?

Are they also unimodal? **Answer:** Not always.

For the circular poset $(1, a, 1, a)$ we get a small dip in the middle:

$$(1, 2, \dots, a, a + 1, a, a + 1, a, a - 1, \dots, 2, 1).$$

Nicer answer: Almost always.

We know the fence poset is unimodal if one of these apply:

What about the rank polynomials of circular fence posets?

Are they also unimodal? **Answer:** Not always.

For the circular poset $(1, a, 1, a)$ we get a small dip in the middle:

$$(1, 2, \dots, a, a + 1, a, a + 1, a, a - 1, \dots, 2, 1).$$

Nicer answer: Almost always.

We know the fence poset is unimodal if one of these apply:

- We have an odd number of nodes.

What about the rank polynomials of circular fence posets?

Are they also unimodal? **Answer:** Not always.

For the circular poset $(1, a, 1, a)$ we get a small dip in the middle:

$$(1, 2, \dots, a, a + 1, a, a + 1, a, a - 1, \dots, 2, 1).$$

Nicer answer: Almost always.

We know the fence poset is unimodal if one of these apply:

- We have an odd number of nodes.
- There are two consecutive parts that are larger than 1.

What about the rank polynomials of circular fence posets?

Are they also unimodal? **Answer:** Not always.

For the circular poset $(1, a, 1, a)$ we get a small dip in the middle:

$$(1, 2, \dots, a, a + 1, a, a + 1, a, a - 1, \dots, 2, 1).$$

Nicer answer: Almost always.

We know the fence poset is unimodal if one of these apply:

- We have an odd number of nodes.
- There are two consecutive parts that are larger than 1.
- There are three consecutive parts $k, 1, l$ with $|k - l| > 1$.

What about the rank polynomials of circular fence posets?

Are they also unimodal? **Answer:** Not always.

For the circular poset $(1, a, 1, a)$ we get a small dip in the middle:

$$(1, 2, \dots, a, a + 1, a, a + 1, a, a - 1, \dots, 2, 1).$$

Nicer answer: Almost always.

We know the fence poset is unimodal if one of these apply:

- We have an odd number of nodes.
- There are two consecutive parts that are larger than 1.
- There are three consecutive parts $k, 1, l$ with $|k - l| > 1$.

We also know that if there is a problem with unimodality, it only happens in the middle.

What about the rank polynomials of circular fence posets?

Are they also unimodal? **Answer:** Not always.

For the circular poset $(1, a, 1, a)$ we get a small dip in the middle:

$$(1, 2, \dots, a, a + 1, a, a + 1, a, a - 1, \dots, 2, 1).$$

Nicer answer: Almost always.


Conjecture (Kantarci Oğuz, Ravichandran, 2021)

For any $\alpha \neq (1, k, 1, k)$ or $(k, 1, k, 1)$ for some k , the rank sequence $\bar{R}(\alpha; q)$ is unimodal.

Another Perspective

We can also see fences as intervals in the Young's lattice.

Young's Lattice is the lattice of Ferrers diagrams of Partitions ordered by inclusion.




(Image from Wikipedia, created by David Eppstein)

For any partition, we can look at the generating function of the partitions that lay under it.

$$G(\lambda; q) := \sum_{\mu \subset \lambda} q^{|\mu|}$$


$$G\left(\begin{array}{|c|} \hline \square \\ \hline \square \\ \hline \end{array}; q\right) = q^3 + 2q^2 + q + 1$$

$$G\left(\begin{array}{|c|} \hline \square \\ \hline \square \\ \hline \square \\ \hline \end{array}; q\right) = q^4 + 2q^3 + 2q^2 + q + 1$$



For any partition, we can look at the generating function of the partitions that lay under it.

$$G(\lambda; q) := \sum_{\mu \subset \lambda} q^{|\mu|}$$




$$G\left(\begin{array}{|c|} \hline \square \\ \hline \square \\ \hline \end{array}; q\right) = q^3 + 2q^2 + q + 1$$

$$G\left(\begin{array}{|c|} \hline \square \\ \square \\ \hline \square \\ \hline \end{array}; q\right) = q^4 + 2q^3 + 2q^2 + q + 1$$

We can also look at the interval between two partitions.

For any partition, we can look at the generating function of the partitions that lay under it.

$$G(\lambda; q) := \sum_{\mu \subset \lambda} q^{|\mu|}$$



$$G\left(\begin{array}{|c|} \hline \square \\ \hline \square \\ \hline \end{array}; q\right) = q^3 + 2q^2 + q + 1$$

$$G\left(\begin{array}{|c|} \hline \square \\ \hline \square \\ \hline \square \\ \hline \end{array}; q\right) = q^4 + 2q^3 + 2q^2 + q + 1$$

We can also look at the interval between two partitions.


$$G(\lambda/\nu; q) := \sum_{\nu \subset \mu \subset \lambda} q^{|\mu| - |\nu|}$$

$$G\left(\begin{array}{|c|} \hline \square \\ \hline \square \\ \hline \end{array} / \begin{array}{|c|} \hline \square \\ \hline \end{array}; q\right) = q^2 + 2q + 1$$

Unimodality of these polynomials were considered by Stanton in 1990⁵.


⁵Stanton, “Unimodality and Young’s lattice” .

Unimodality of these polynomials were considered by Stanton in 1990⁵. Note that taking the transpose does not change the polynomial we get, so we can think up to transpose.



⁵Stanton, “Unimodality and Young’s lattice”.

Unimodality of these polynomials were considered by Stanton in 1990⁵. Note that taking the transpose does not change the polynomial we get, so we can think up to transpose.



Conjecture (Stanton,1990)

The polynomials corresponding to self-dual partitions are unimodal.

⁵Stanton, “Unimodality and Young’s lattice”.

The counter examples mainly occur in the case where we have 4 parts, where we only get a dip in the middle.

The counter examples mainly occur in the case where we have 4 parts, where we only get a dip in the middle.


TABLE I

Partition	i	Values	Partition	i	Values
8 8 4 4	15	31 30 31	11 11 6 6	21	67 66 67
10 9 4 4	17	46 45 46	14 13 4 4	21	76 75 76
10 10 4 4	17	46 45 46	16 12 4 4	23	91 90 91
12 10 4 4	19	61 60 61	14 14 4 4	21	76 75 76
12 11 4 4	19	61 60 61	12 12 8 4	23	81 80 81
12 12 4 4	19	61 60 61	12 10 8 6	23	82 81 82
14 11 4 4	21	76 75 76	8 8 8 6 4 2	23	141 140 141
11 11 6 5	21	67 66 67	8 8 6 6 4 4	23	144 143 144
14 12 4 4	21	76 75 76			

(Table from "Unimodality and Young's Lattice", Stanton)


Given a fence, we can see it as a difference of two partitions α/ν .

Example $((2, 1, 1, 3) \rightarrow (4, 4, 4, 4, 3)/(3, 3, 3, 2))$



Given a fence, we can see it as a difference of two partitions α/ν .


Example $((2, 1, 1, 3) \rightarrow (4, 4, 4, 4, 3)/(3, 3, 3, 2))$



Note that the ideals of the fence coincide with the partitions that lie between α and ν , so $G(\lambda/\nu)$ agrees with the rank polynomial.

Given a fence, we can see it as a difference of two partitions α/ν .

Example $((2, 1, 1, 3) \rightarrow (4, 4, 4, 4, 3)/(3, 3, 3, 2))$




Note that the ideals of the fence coincide with the partitions that lie between α and ν , so $G(\lambda/\nu)$ agrees with the rank polynomial.

Rank polynomials actually correspond to a special class of differences called *ribbon diagrams*, where we have no 2×2 box.

Given a fence, we can see it as a difference of two partitions α/ν .

Example $((2, 1, 1, 3) \rightarrow (4, 4, 4, 4, 3)/(3, 3, 3, 2))$








Note that the ideals of the fence coincide with the partitions that lie between α and ν , so $G(\lambda/\nu)$ agrees with the rank polynomial.

Rank polynomials actually correspond to a special class of differences called *ribbon diagrams*, where we have no 2×2 box.

Polynomials corresponding to ribbon diagrams are unimodal.

Thank you for listening!

Further Reading

-  Kantarcı Oğuz, E. & Ravichandran, M. Rank Polynomials of Fence Posets are Unimodal. (2021)
-  Morier-Genoud, S. & Ovsienko, V. q -deformed rationals and q -continued fractions. *Forum Math. Sigma*. **8** pp. Paper No. e13, 55 (2020).
-  McConville, T., Sagan, B. & Smyth, C. On a rank-unimodality conjecture of Morier-Genoud and Ovsienko. *Discrete Math.* **344** pp. 13 (2021).
-  Elizalde, S. & Sagan, B. Partial rank symmetry of distributive lattices for fences. (2022)
-  Stanton, D. Unimodality and Young's lattice. *J. Comb. Theory, Ser. A*. **54**, 41-53 (1990)