

Binary Quadratic Forms as Dessins

A. Muhammed Uludağ, Ayberk Zeytin, Merve Durmuş

October 8, 2012

Abstract

We show that the class of every primitive indefinite binary quadratic form is naturally represented by an infinite graph (named *çark*) with a unique cycle embedded on a conformal annulus. This cycle is called the spine of the *çark*. Every *çark* is an infinite dessin and determines an annular uniformization of the modular curve. Every choice of a base edge of a fixed *çark* specifies an indefinite binary quadratic form in the class represented by the *çark*. The proper automorphism group of a form is identified with the fundamental group of its *çark*. Reduced forms in the class represented by a *çark* correspond to some distinguished base edges of its spine. The reduction algorithm of Gauss is the process of moving the base edge in the direction of the spine of the *çark*. Ambiguous and reciprocal classes are represented by *çarks* with symmetries. Periodic *çarks* represent classes of non-primitive forms.

1 Introduction.

The Euclidean algorithm is the process of comparison of commensurable magnitudes and the modular group $\mathrm{PSL}_2(\mathbb{Z})$ is the encoding of this algorithm. Since the intellect is ultimately about comparison of magnitudes, it should come as no surprise that the modular group manifests itself in diverse contexts through its action on mathematical objects, no matter what our level of abstraction is. Together with its action on plane lattices, among all manifestations of $\mathrm{PSL}_2(\mathbb{Z})$ the following four classical actions are of fundamental nature:

1. its left-action on the infinite trivalent plane tree,
2. its left action on the upper half plane \mathcal{H} by Möbius transformations,
3. its right-action on the binary quadratic forms of Gauss, and
4. its left-conjugation action on itself.

In this paper we clarify some connections among these four actions. The first action is not free on the set of neither edges nor vertices of the tree in question. In order to make it free on the set of edges, we add the midpoints as extra vertices thereby doubling the set of edges and call the resulting infinite bipartite tree the *Farey tree* \mathcal{F} . In fact it is possible to construct \mathcal{F} in a purely combinatorial manner which we will carry out in Section 2, starting from a pair of elliptic generators of

$\mathrm{PSL}_2(\mathbb{Z})$. Now since $\mathrm{PSL}_2(\mathbb{Z})$ acts on \mathcal{F} by automorphisms; freely on the set of edges of \mathcal{F} , so does any subgroup Γ of $\mathrm{PSL}_2(\mathbb{Z})$, and by our definition a *modular graph*¹ is simply a quotient graph $\Gamma \backslash \mathcal{F}$. This is almost the same thing as a trivalent ribbon graph, except that we consider the midpoints as extra 2-valent vertices and pending edges are allowed. Modular graphs parametrize subgroups of the modular group up to conjugacy and modular graphs with a base edge classify subgroups of the modular group.


The second action is compatible with the first one in the following sense: The tree $\mathcal{F}_{top} \subset \mathcal{H}$ which is built as the $\mathrm{PSL}_2(\mathbb{Z})$ -orbit of the arc connecting two elliptic points on the boundary of the standard fundamental domain, is a topological realization of the Farey tree \mathcal{F} . Consequently, $\Gamma \backslash \mathcal{F}_{top} \subset \Gamma \backslash \mathcal{H}$ is a topological realization of the graph $\Gamma \backslash \mathcal{F}$, as a graph embedded in the Riemann orbifold $\Gamma \backslash \mathcal{H}$. This latter is a surface (i.e. it has no orbifold points) in case Γ contains no elliptics but always has punctures due to the parabolic elements of Γ , or it has some boundary components. These punctures are in one-to-one correspondence with the left-turn circuits in $\Gamma \backslash \mathcal{F}$. Widening these punctures gives a deformation retract of the ambient orbifold to the graph, in particular the upper half plane \mathcal{H} retracts to the Farey tree \mathcal{F}_{top} . To recover the (topological) orbifold from the modular graph one glues punctured discs along the left-turn paths of the graph.

In case Γ is of finite index in $\mathrm{PSL}_2(\mathbb{Z})$, the quotient $\Gamma \backslash \mathcal{H}$ is an algebraic curve (in fact a stack) which can be defined over a number field since it is a finite covering of the modular curve $\mathcal{M} = \mathrm{PSL}_2(\mathbb{Z}) \backslash \mathcal{H}$. According to Belyı̄'s theorem, [2], any arithmetic surface can be defined this way, implying in particular that the action of the absolute Galois group defined on the set of finite coverings $\{\Gamma \backslash \mathcal{H} \rightarrow \mathcal{M}\}$ is faithful. But these coverings are equivalently described by the graphs $\Gamma \backslash \mathcal{F}$. This striking correspondence between combinatorics and arithmetic led Grothendieck and his followers to study *dessins* (=modular graphs) from the point of view of the action of the absolute Galois group, see [16]. However, explicit computations of covering maps $\Gamma \backslash \mathcal{H} \rightarrow \mathcal{M}$ required by this approach turned out to be forbiddingly hard if one wants to go beyond some basic cases and only a few uniform theorems could be obtained. In fact, *dessins* are more general graphs that correspond to finite coverings of the thrice punctured sphere, which is equivalent to a subsystem of coverings of \mathcal{M} since $\mathbb{P}^1 \setminus \{0, 1, \infty\}$ is a degree-6 covering of \mathcal{M} (because the congruence subgroup of level 2 is an index-6 subgroup in the modular group). So modular graphs include *dessins* and hence there is no loss of generality in considering modular graphs.


Turning back to our four actions, in the third one the modular group acts on the set of binary quadratic forms by acting on the pair of variables in the well-known manner. Orbits of this action are called *classes* and forms in the same class are said to be *equivalent*. This action was studied systematically by Gauss. Here we are interested in the action on *indefinite* forms. This action always has a cyclic stabilizer

¹Adding to the long list of names and equivalent/dual notions with various nuisances: cyclic trivalent graphs, cuboid tree diagrams, Jacobi diagrams, trivalent ribbon graphs, triangulations; more generally, maps, ribbon graphs, fat graphs, *dessins*, polygonal decompositions, lozenge tilings, etc.

group, which is called the proper automorphism group of the form and denoted $\langle M_f \rangle$. Indefinite binary quadratic forms represent ideal classes in the quadratic number field having the same discriminant as the form and hence are tightly related to real quadratic number fields [5]. We provide a succinct introduction to binary quadratic forms later in the paper.



(a) A dessin (linienzug of Klein) from 1879 [14]



(b) A çark in its ambient annulus

The correspondence between forms and dessins is elementary and can be described briefly as follows: to an indefinite binary quadratic form f we associate its proper automorphism group $\langle M_f \rangle$ and to $\langle M_f \rangle$ we associate the infinite graph $\langle M_f \rangle \backslash \mathcal{F}$, which is called a *çark*². Via the topological realization of \mathcal{F} , this is a graph embedded in the annulus $\langle M_f \rangle \backslash \mathcal{H}$. Çarks are infinite “transcendental” graphs whereas the dessins literature consider only finite graphs. (“transcendental” since they correspond to non-algebraic extensions of the function field of the modular curve). This transcendence implies that çarks go undetected in the algebraic fundamental group approach, nevertheless we shall see that this does not keep them away from being arithmetic objects.

Equivalent forms have conjugate stabilizers (automorphism groups) and conjugate subgroups have isomorphic quotient graphs. It turns out that the set of classes is exactly the set of orbits of hyperbolic elements of $\mathrm{PSL}_2(\mathbb{Z})$ under the fourth (conjugation) action in our list. This set of orbits can be identified with the set of bracelet diagrams with beads of two colors.

As we shall see later in the paper, çarks provide a very nice reformulation of various concepts pertaining to indefinite binary quadratic forms, such as reduced forms and the reduction algorithm, ambiguous forms, reciprocal forms, the Markoff value of a form, etc. In fact, çarks are nothing but \mathbb{Z} -quotients of periodic rivers of Conway [6]. However, it must be admitted that we could contribute nothing to the classical and deep questions concerning the behavior of class numbers and the structure of class groups. For us the importance of this elementary correspondence between çarks and forms lies in that it suggests a concrete way to consider modular graphs as arithmetic objects viz. Gauss’ binary quadratic forms, as it was much solicited by Grothendieck’s dessins school.

²Turkish *çark* (pronounced as “chark”) is borrowed from Persian, and it has a common etymology with Indian *chakra*, Greek *kuklos* and English *wheel*.

To conclude, let us rephrase our main result: we show that the class of every primitive indefinite binary quadratic form is not simply a set but it has the extra structure of an infinite graph, namely a çark, such that the forms in the class are identified with the edges of the graph. This graph admits a topological realization as a subset of an annulus and explains very well some known phenomena around Gauss' reduction theory of indefinite binary quadratic forms. Various properties of forms and their classes are manifested in a natural way on the çark.

2 Farey tree and modular graphs

It is well known that, see for instance [4, Excercise 2.9.1], the two elliptic transformations $S(z) = -1/z$ and $R(z) = (z-1)/z$, respectively of orders 2 and 3, generate a group of Möbius transformations which is isomorphic to the projective group of two by two integral matrices having determinant 1, the modular group $\mathrm{PSL}_2(\mathbb{Z})$. It is also well-known that $\mathrm{PSL}_2(\mathbb{Z}) \cong \langle S \rangle * \langle R \rangle = \mathbb{Z}/2\mathbb{Z} * \mathbb{Z}/3\mathbb{Z}$. Let us now consider the graph \mathcal{F} (so called Farey tree), given by the following data:

$$\begin{aligned} E(\mathcal{F}) &= \{\{W\} : W \in \mathrm{PSL}_2(\mathbb{Z})\} \\ V(\mathcal{F}) &= V_{\otimes}(\mathcal{F}) \sqcup V_{\bullet}(\mathcal{F}); \end{aligned}$$

where

$$\begin{aligned} V_{\otimes}(\mathcal{F}) &= \{\{W, WS\} : W \in \mathrm{PSL}_2(\mathbb{Z})\}, \text{ and} \\ V_{\bullet}(\mathcal{F}) &= \{\{W, WR, WR^2\} : W \in \mathrm{PSL}_2(\mathbb{Z})\}. \end{aligned}$$

There is an edge between a vertex $v = \{W, WS\} \in V_{\otimes}(\mathcal{F})$ and another vertex $v' = \{W', W'R, W'R^2\}$ if and only if $\{W, WS\} \cap \{W', W'R, W'R^2\} \neq \emptyset$ and there are no other edges. Thus the edge connecting v and v' is $v \cap v'$, if this intersection is non-empty. Observe that by construction the graph is bipartite. The edges incident to the vertex $\{W, WR, WR^2\} \in V_{\bullet}$ are $\{W\}, \{WR\}, \{WR^2\}$, and these edges inherit a natural cyclic ordering from the vertex. Thus the Farey tree \mathcal{F} is an infinite bipartite ribbon graph.

The group $\mathrm{PSL}_2(\mathbb{Z})$ acts on \mathcal{F} from the left, by ribbon graph automorphisms, where $M \in \mathrm{PSL}_2(\mathbb{Z})$ acts by

$$\begin{aligned} \{W\} \in E(\mathcal{F}) &\rightarrow \{MW\} \in E(\mathcal{F}) \\ \{W, WS\} \in V_{\otimes}(\mathcal{F}) &\rightarrow \{MW, MWS\} \in V_{\otimes}(\mathcal{F}) \\ \{W, WR, WR^2\} \in V_{\bullet}(\mathcal{F}) &\rightarrow \{MW, MWR, MWR^2\} \in V_{\bullet}(\mathcal{F}) \end{aligned}$$

Notice that the action on the set of edges is nothing but the left-regular action of $\mathrm{PSL}_2(\mathbb{Z})$ on itself and therefore is free. On the other hand the action is not free on the set of vertices: The vertex $\{W, WS\}$ is fixed by the order-2 subgroup generated by $M = WSW^{-1}$, and the vertex $\{W, WR, WR^2\}$ is fixed by the order-3 subgroup generated by $M = WRW^{-1}$.

Let Γ be any subgroup of $\mathrm{PSL}_2(\mathbb{Z})$. Then Γ acts on \mathcal{F} from the left and to Γ we associate a quotient graph $\Gamma \backslash \mathcal{F}$ as follows:

$$E(\Gamma \backslash \mathcal{F}) = \{\Gamma \cdot \{W\} : W \in \mathrm{PSL}_2(\mathbb{Z})\}$$

$$V(\Gamma \backslash \mathcal{F}) = V_{\otimes}(\mathcal{F}/\Gamma) \cup V_{\bullet}(\mathcal{F}/\Gamma);$$

where

$$\begin{aligned} V_{\otimes}(\Gamma \backslash \mathcal{F}) &= \{\Gamma \cdot \{W, WS\} : W \in \mathrm{PSL}_2(\mathbb{Z})\}, \text{ and} \\ V_{\bullet}(\Gamma \backslash \mathcal{F}) &= \{\Gamma \cdot \{W, WR, WR^2\} : W \in \mathrm{PSL}_2(\mathbb{Z})\}. \end{aligned}$$

It is easy to see that the incidence relation induced from the Farey tree gives a well-defined incidence relation and gives us the graph which we call a *modular graph*. Thus the edge connecting the vertices $v = \Gamma \cdot \{W, WS\}$ and $v' = \Gamma \cdot \{W', W'R, W'R^2\}$ is the intersection $v \cap v'$, which is of the form $\Gamma \cdot \{M\}$ if non-empty. There are no other edges. Observe that by construction the graph is bipartite. The edges incident to the vertex $\Gamma \cdot \{W, WR, WR^2\}$ are $\Gamma \cdot \{W\}$, $\Gamma \cdot \{WR\}$, $\Gamma \cdot \{WR^2\}$, and these edges inherit a natural cyclic ordering from the vertex. Thus the Farey tree \mathcal{F} is an infinite bipartite ribbon graph³. In general $\Gamma \backslash \mathcal{F}$ is a bipartite ribbon graph possibly with pending vertices that corresponds to the conjugacy classes of elliptic elements that Γ contains. Conversely, any connected bipartite ribbon graph G , with $V(G) = V_{\otimes}(G) \sqcup V_{\bullet}(G)$, such that every \otimes -vertex is of degree 1 or 2 and every \bullet -vertex is of degree 1 or 3, is modular since the universal covering of G is isomorphic to \mathcal{F} . As in [18], it takes a little effort to define the fundamental group of $\Gamma \backslash \mathcal{F}$ so that there is a canonical isomorphism $\pi_1(\Gamma \backslash \mathcal{F}, \Gamma \cdot \{I\}) \simeq \Gamma < \mathrm{PSL}_2(\mathbb{Z})$, with the canonical choice of $\Gamma \cdot \{I\}$ as a base edge. In general, subgroups Γ of the modular group (or equivalently the fundamental groups $\pi_1(\Gamma \backslash \mathcal{F})$) are free products of copies of \mathbb{Z} , $\mathbb{Z}/2\mathbb{Z}$ and $\mathbb{Z}/3\mathbb{Z}$, see [15]. Note that two distinct isomorphic subgroups Γ_1 , Γ_2 of the modular group may give rise to non-isomorphic ribbon graphs $\Gamma_1 \backslash \mathcal{F}$ and $\Gamma_2 \backslash \mathcal{F}$. In other words, the fundamental group does not characterize the graph. Another basic invariant of $\Gamma \backslash \mathcal{F}$ is its genus, which is defined to be the genus of the surface constructed by gluing discs along right-turn paths. This genus is the same as the genus of the algebraic curve \mathcal{H}/Γ .

The set of edges of $\Gamma \backslash \mathcal{F}$ is identified with the set of right-cosets of Γ , so that the graph $\Gamma \backslash \mathcal{F}$ has $[\Gamma : \mathrm{PSL}_2(\mathbb{Z})]$ many edges. In case Γ is a finite index subgroup, the graph $\Gamma \backslash \mathcal{F}$ is finite. In case $\Gamma = \mathrm{PSL}_2(\mathbb{Z})$, the quotient graph $\mathrm{PSL}_2(\mathbb{Z}) \backslash \mathcal{F}$ is a graph with one edge that looks like as follows:

$$\begin{array}{ccc} \mathrm{PSL}_2(\mathbb{Z})\{I, S\} & \mathrm{PSL}_2(\mathbb{Z})\{I\} & \mathrm{PSL}_2(\mathbb{Z})\{I, R, R^2\} \\ \otimes & \text{---} & \bullet \end{array}$$

Figure 1: The modular arc.

We call this graph the *modular arc*. It is a graph whose fundamental group is $\mathrm{PSL}_2(\mathbb{Z})$ and whose universal cover is the Farey tree \mathcal{F} . In other words modular graphs are coverings of the modular arc. If we consider the action of the modular group on the topological realization \mathcal{F}_{top} of \mathcal{F} mentioned in the introduction, the topological realization of $\mathrm{PSL}_2(\mathbb{Z}) \backslash \mathcal{F}$ is the arc $\mathrm{PSL}_2(\mathbb{Z}) \backslash \mathcal{F}_{top}$ in the modular curve connecting two elliptic points.

Every modular graph $\Gamma \backslash \mathcal{F}$ has a canonical ‘‘analytical’’ realization $\Gamma \backslash \mathcal{F}_{top}$ on the Riemann surface $\Gamma \backslash \mathcal{H}$ with edges being geodesic arcs. Equivalently, these edges

³The ribbon graph structure around vertices of degree 2 is trivial.




Figure 2: (Right) The fundamental region for the modular curve in the upper half plane model. (Left) The modular curve. Note that there are two triangles, the second is on the back of the page, glued to this one. SAGDAKI SEKILIN KOSE NOKTALARINI BELIRTILECEK, SOLDAGINDE OLDUGU GIBI

are lifts of the modular arc by $\Gamma \backslash \mathcal{H} \rightarrow \mathrm{PSL}_2(\mathbb{Z}) \backslash \mathcal{H}$. If instead we lift the geodesic arc connecting the \otimes - elliptic point to the cusp to the surface $\Gamma \backslash \mathcal{H}$, then we obtain another graph on the surface, which is called an *ideal triangulation*. Lifting the remaining geodesic arc gives rise to yet another type of graph, called a *lozenge tiling*. So there is a triality, not just duality, of these graphs.




Figure 3: Triality of graphs

In topology, there is a well-known correspondence between subgroups of the fundamental group of a space and the coverings of that space. The following two results are orbifold (or “stacky”) analogues of this correspondence for coverings of the modular curve, stated in terms of graphs. For more details on fundamental groups and covering theory of graphs see [18].

Proposition 2.1. *If Γ_1 and Γ_2 are conjugate subgroups of $\mathrm{PSL}_2(\mathbb{Z})$, then the graphs $\Gamma_1 \backslash \mathcal{F}$ and $\Gamma_2 \backslash \mathcal{F}$ are isomorphic as ribbon graphs. Hence there is a 1-1 correspondence between modular graphs and conjugacy classes of subgroups of the modular group.*

Proof. Let $\Gamma_2 = M\Gamma_1 M^{-1}$. The desired isomorphism is then the map

$$\varphi : \Gamma_1 \cdot \{W\} \in E(\Gamma_1 \backslash \mathcal{F}) \rightarrow \Gamma_2 \cdot \{MW\} \in E(\Gamma_2 \backslash \mathcal{F}).$$

Note that one has $\varphi(\Gamma_1 \cdot \{I\}) = \Gamma_2 \cdot \{M\}$. Suppose now that $\varphi : E(\Gamma_1 \backslash \mathcal{F}) \rightarrow E(\Gamma_2 \backslash \mathcal{F})$

is a ribbon graph isomorphism and let $\varphi(\Gamma_1 \cdot \{I\}) = \Gamma_2 \cdot \{M\}$. This induce an isomorphism of fundamental groups

$$\varphi_* : \pi_1(\Gamma_1 \backslash \mathcal{F}, \Gamma_1 \cdot \{I\}) \simeq \pi_1(\Gamma_2 \backslash \mathcal{F}, \Gamma_2 \cdot \{M\})$$

Since φ is a ribbon graph isomorphism, these two groups are also isomorphic as subgroups of the modular group. The former group is canonically isomorphic to Γ_1 whereas the latter group is canonically isomorphic to

$$M^{-1}\pi_1(\Gamma_2 \backslash \mathcal{F}, \Gamma_2 \cdot \{I\})M \simeq M^{-1}\Gamma_2M \quad \square$$

Therefore modular graphs parametrize conjugacy classes of subgroups of the modular group, whereas the edges of a modular graph parametrize subgroups in the conjugacy class represented by the modular graph, and we get:

Theorem 2.2. *There is a 1-1 correspondence between modular graphs with a base edge (G, e) (modulo ribbon graph isomorphisms of pairs (G, e)) and subgroups of the modular group.*

3 Çarks

A *çark* is a modular graph which is of the form $\mathcal{C}_M := \langle M \rangle \backslash \mathcal{F}$ where M is a hyperbolic element of the modular group. One has

$$\pi_1(\langle M \rangle \backslash \mathcal{F}) = \langle M \rangle \simeq \mathbb{Z},$$

so the çark $\langle W \rangle \backslash \mathcal{F}$ is a graph with only one circuit, which we call the *spine* of the çark. Every çark has a canonical realization as a graph $\langle M \rangle \backslash \mathcal{F}_{top}$ embedded in the surface $\langle M \rangle \backslash \mathcal{H}$, which is an annulus since M is hyperbolic. In fact $\langle M \rangle \backslash \mathcal{H}$ is the annular uniformization of the modular curve \mathcal{M} corresponding to $M \in \pi_1(\mathcal{M})$. Again by hyperbolicity of M , this graph will have infinite ‘‘Farey branches’’ attached to the spine in the direction of both of the boundary components of the annulus⁴. By Proposition 2.1 the graphs \mathcal{C}_M and $\mathcal{C}_{XMX^{-1}}$ are isomorphic for every element X of the modular group and by Theorem 2.2 we deduce the following result, see [7]:

Corollary 3.1. *There are one-to-one correspondences between*

- i. çarks and conjugacy classes of subgroups of the modular group generated by a single hyperbolic element, and*
- ii. çarks with a base edge and subgroups of the modular group generated by a single hyperbolic element.*

⁴If M is parabolic, then $\langle W \rangle \backslash \mathcal{F}$ has Farey branches attached to the spine in only one direction, and its topological realization $\langle M \rangle \backslash \mathcal{F}_{top}$ sits on a punctured disc. If M is elliptic, $\langle W \rangle \backslash \mathcal{F}$ is a tree with a pending edge which abut at a vertex of type \otimes when M is of order 2 and of type \bullet when M is of order 3. Its topological realization $\langle M \rangle \backslash \mathcal{F}_{top}$ sits on a disc with an orbifold point.




Figure 4: The çark $\mathcal{F}/\langle SR^2SR \rangle$.

Corollary 3.2. *There are one-to-one correspondences between*

- i. hyperbolic elements of the modular group and directed çarks with a base edge, and*
- ii. conjugacy classes of hyperbolic elements of the modular group and directed çarks.*

3.1 Necklaces and Bracelets

Çarks are infinite graphs, and according to our definition each edge of a çark carries a name which is an infinite coset. In fact all the combinatorial information of a çark can be encoded in a finite storage as follows: First remove all \circ -vertices of the çark. Next, turn once around the spine, whenever you meet a \bullet -vertex on which a branch attached by R , cut that branch and tag that \bullet -vertex with a “0”, tagging all the remaining vertices by a “1” similarly cutting the branches attached to them. We obtain a finite graph called a *binary bracelet* which is by definition an equivalence class of binary strings under cyclic permutations (i.e. rotations) and reversals. Conversely, by using the convention $0 \leftrightarrow R$ and $1 \leftrightarrow R^2$ we can reconstruct the çark from its bracelet.

Rotations and reversals generate a finite dihedral group, and a binary bracelet may equivalently be described as an orbit of this action. For $n = 1, 2, \dots, 15$ the number of binary bracelets is

$$2, 3, 4, 6, 8, 13, 18, 30, 46, 78, 126, 224, 380, 687, 1224.$$

This is sequence A000029 (M0563) in OEIS [1]. The number of binary bracelets (çarks) of length n is

$$B(n) = \begin{cases} \frac{1}{2}N(n) + \frac{3}{4}2^{n/2} & \text{if } n \text{ is even} \\ \frac{1}{2}N(n) + \frac{1}{2}2^{(n+1)/2} & \text{if } n \text{ is odd,} \end{cases}$$

where $N(n)$ is the number of binary necklaces of length n . An equivalence class of binary strings under rotations (excluding thus reversals) is called a *binary necklace*,




Figure 5: From çarks to bracelets

also called a *cyclic binary word*. They are thus orbits of words under the action of a cyclic group and they correspond to directed çarks. For $n = 1, 2, \dots, 15$ the number of binary necklaces of length n is

$$N(n) = 2, 3, 4, 6, 8, 14, 20, 36, 60, 108, 188, 352, 632, 1182, 2192,$$

which is sequence A000031 (M0564) in OEIS. The number of necklaces (directed çarks) of length n is given by MacMahon's formula from 1892 (also called Witt's formula) (see [3], [17])

$$N(n) = \frac{1}{n} \sum_{d|n} \varphi(d) 2^{n/d} = \frac{1}{n} \sum_{j=1}^n 2^{\gcd(j,n)}$$

where φ is Euler's totient function.

Aperiodic binary necklaces correspond to primitive directed çarks. For $n = 1, 2, \dots, 15$ the number of aperiodic necklaces of length n is

$$L(n) = 2, 1, 2, 3, 6, 9, 18, 30, 56, 99, 186, 335, 630, 1161, 2182,$$

which is sequence A000031 (M0564) in the database. There is a formula for the number of aperiodic necklaces of length n in terms of Möbius' function μ :

$$L(n) = \frac{1}{n} \sum_{d|n} \mu(d) 2^{n/d} = \frac{1}{n} \sum_{d|n} \mu(n/d) 2^d$$

As we said, binary necklaces (=cyclic binary words=directed çarks) may be viewed as orbits of words under the action of the cyclic group. Choosing an ordering of our letters $\{0, 1\}$ (i.e. $0 < 1$) and imposing the lexicographic ordering of the words, one may choose a minimal representative in each orbit. The minimal representative of a primitive (aperiodic) word is called a *Lyndon word*. They were first studied in connection with the construction of bases for free lie algebras and they appear in numerous contexts. In our case they are

$$0, 1, 01, 001, 011, 0001, 0011, 0111, 00001, 00011, 00101, 00111, 01011, 01111 \dots$$

One can similarly find representatives for aperiodic binary bracelets (=primitive indefinite binary quadratic forms; see below). There are effective algorithms to list all primitive necklaces and bracelets up to a given length (i.e. Duval’s algorithm [8], the algorithm due to Fredricksen, Kessler and Maiorana [9], Sawada’s algorithm [20], etc). Translated into the language of binary quadratic forms, this means that it is possible to single out a unique reduced representative in each class of a primitive indefinite binary quadratic form and that it is possible to effectively enumerate all classes of primitive indefinite binary quadratic forms by specifying those reduced representatives.

To sum up, we may represent primitive çarks by primitive bracelets. In order to shorten this representation further, we may count the number of consecutive 0’s and 1’s and represent çarks as sequences of natural numbers $(n_0, n_1, \dots, n_{2k})^{0,1}$, if we agree that⁵ this sequence encodes a bracelet that starts with a 0 if the exponent is 0 and 1 if the exponent is 1. This representation is directly connected to the “minus” continued fractions (see Zagier [22]).

A primitive word may have two types of symmetries: invariance under the swap of symbols $0 \leftrightarrow 1$ and invariance under reversal, i.e. palindromic symmetry. The first symmetry corresponds to ambiguous binary quadratic forms and the second symmetry corresponds to reciprocal binary quadratic forms, as we shall see. The swap of symbols $0 \leftrightarrow 1$ corresponds to inversion in the class group.

3.2 Çark Invariants

There are several natural invariants associated to a çark \mathcal{C} . The first invariant that comes to mind is the combinatorial length $l_c(\mathcal{C})$ of its spine. A hyperbolic invariant of a çark is the metric length $l_h(\mathcal{C})$ of the closed geodesic in the annular surface under its hyperbolic metric induced by the çark. A conformal invariant of a çark is the modulus $m(\mathcal{C})$ of the associated annulus. Finally, the discriminant $\Delta(\mathcal{C})$ of the associated form and the trace $\tau(\mathcal{C})$ of the associated matrix are two arithmetic invariants with $\Delta = \tau^2 - 4$. One has

$$l_h(\mathcal{C}) = 2 \operatorname{arccosh}(\tau/2), \quad m(\mathcal{C}) = \exp\left(\frac{\pi^2}{\log|\frac{\tau \pm \sqrt{\Delta}}{2}|}\right)$$

The modulus is found as follows: Any hyperbolic element $M \in PSL_2(\mathbb{R})$ is conjugate to an element of the form

$$N := XMX^{-1} = \begin{bmatrix} \alpha & 0 \\ 0 & 1/\alpha \end{bmatrix}$$

where α is the multiplier of M . Since the trace is invariant under conjugation, one has $\tau := \operatorname{tr}(M) = \alpha + 1/\alpha \implies \alpha^2 - \tau\alpha + 1 = 0 \implies \alpha = \frac{\tau \pm \sqrt{\tau^2 - 4}}{2}$.

Now N acts by Möbius transformation $z \mapsto \alpha^2 z$, and the quotient map is $f(z) = z^{2\pi i / \log \alpha^2}$ with the annulus $f(\mathcal{H}) = \{z : e^{-2\pi^2 / \log \alpha^2} < |z| < 1\}$ as its image. Hence the modulus of the ambient annulus of the çark is $e^{2\pi^2 / \log \alpha^2} = e^{\pi^2 / \log |\alpha|}$.

⁵Note that a Lyndon word always start with a 0 and ends with a 1.

4 Binary Quadratic Forms and Çarks

A *binary quadratic form* is a homogeneous function of degree two in two variables $f(x, y) = Ax^2 + Bxy + Cy^2$ (denoted $f = (A, B, C)$ or in the matrix form:

$$W_f = \begin{pmatrix} A & B/2 \\ B/2 & C \end{pmatrix} \quad (4.1)$$

so that $f(x, y) = (x, y)W_f(x, y)^t$. If the coefficients A, B, C are integers the form is called *integral* with discriminant $\Delta(f) = B^2 - 4AC$. If f is integral and $\gcd(A, B, C) = 1$ then f is called *primitive*. Following Gauss we will call a form $f = (A, B, C)$ *ambiguous* if $B = kA$ for some $k \in \mathbb{Z}$. Finally a form $f = (A, B, C)$ will be referred to as *reciprocal* whenever $C = -A$, [19].

Note that $\Delta(f) = -4 \det(W_f)$. Given a symmetric two by two matrix we write f_W to denote the binary quadratic form associated to W . Recall that a form f is called

- positive definite if and only if $\Delta(f) < 0$ and $A > 0$,
- negative definite if and only if $\Delta(f) < 0$ and $0 > A$,
- indefinite if and only if $\Delta(f) > 0$.

The group $\text{PGL}_2(\mathbb{Z})$ acts on the set of all integral binary quadratic forms by

$$\begin{aligned} \text{Forms} \times \text{PGL}(2, \mathbb{Z}) &\rightarrow \text{Forms} \\ (f, U) &\mapsto U \cdot f := \det U f(U(x, y)) \\ &= (\det U)(x, y)U^t W_f U(x, y)^t \end{aligned}$$

We call two binary quadratic forms *equivalent* if they belong to the same $\text{PGL}_2(\mathbb{Z})$ orbit under the above action, under which discriminant is invariant. In particular, two binary quadratic forms are said to be *properly equivalent* if they belong to the same $\text{PSL}_2(\mathbb{Z})$ orbit. Let us denote the $\text{PSL}_2(\mathbb{Z})$ -orbit (or the equivalence class) of f by $[f]$. The stabilizer of f is called its *automorphism group* and denoted by $\text{Aut}(f)$. If we restrict the action to $\text{PSL}_2(\mathbb{Z})$ then the stabilizer is called the *proper automorphism group* of f and denoted by $\text{Aut}_+(f)$. Elements of $\text{Aut}(f)$, (respectively $\text{Aut}_+(f)$) are called automorphisms (respectively proper automorphisms) of f . For a positive definite binary quadratic form f , the group $\text{Aut}(f)$ is trivial unless $\Delta(f) = -3$ or -4 ; $\text{Aut}(f) \simeq \mathbb{Z}/4\mathbb{Z}$ if $\Delta(f) = -4$ and $\text{Aut}(f) \simeq \mathbb{Z}/6\mathbb{Z}$ in case $\Delta(f) = -3$, [4, p.29]. On the other hand, for an indefinite binary quadratic form one has $\text{Aut}(f) \simeq \mathbb{Z}/2\mathbb{Z} \oplus \mathbb{Z}$ and $\text{Aut}_+(f) \simeq \mathbb{Z}$.

Given an indefinite binary quadratic form $f = (A, B, C)$ a generator of its proper automorphism group will be called its *fundamental automorphism*. Note that there are two fundamental automorphisms, one being M_f , the other being its inverse, M_f^{-1} . Every integral solution (α, β) of Pell's equation:

$$X^2 + \Delta(f)Y^2 = +4???????????????? \quad (4.2)$$

corresponds to an automorphism of f given by the matrix:

$$\begin{pmatrix} \frac{x-By}{2} & -Cy \\ Ay & \frac{x+yB}{2} \end{pmatrix}.$$

The fundamental automorphism is the one having minimal β [4, Proposition 6.12.7].

Conversely, to any given hyperbolic element, say $M = \begin{pmatrix} p & q \\ r & s \end{pmatrix} \in \text{PSL}_2(\mathbb{Z})$ let us associate the following binary quadratic form:

$$f_M = \frac{\text{sgn}(p+s)}{\text{gcd}(q, s-p, r)}(r, s-p, -q) \quad (4.3)$$

Observe first that $M \rightarrow f_M$ is well-defined and that its image is always primitive and indefinite. At this point let us state a direct consequence of Theorem ??:

Corollary 4.1. *The maps $\langle M \rangle \backslash \mathcal{F} \leftrightarrow M \rightarrow f_M$ defines a surjection from the set of oriented çarks with a base edge to primitive indefinite binary quadratic forms.*

Proof. We saw that an oriented çark with a base edge determines a hyperbolic element of $\text{PSL}_2(\mathbb{Z})$. And this element in turn determines an indefinite binary quadratic form via $M \rightarrow f_M$. Conversely, given a primitive indefinite binary quadratic form $f = (A, B, C)$ to find $\beta \in \mathbb{Z}$ such that the matrix

$$\begin{pmatrix} \beta & A \\ -C & B + \beta \end{pmatrix} \in \text{PSL}_2(\mathbb{Z})$$

we look at solutions (x, y) of Pell's equation $X^2 + \Delta(f)Y^2 = 4$. Using any such y we construct the hyperbolic element:

$$M_f = \begin{pmatrix} \beta & yC \\ yA & yB + \beta \end{pmatrix},$$

where $\beta = \frac{-yB \pm x}{2}$. Both choices of the sign produces a matrix which maps onto f . In fact, the two matrices are inverses of each other in $\text{PSL}_2(\mathbb{Z})$. \square

Example 4.1. Consider the form $(1, 7, -1)$. It has discriminant 53. The pair $(51, 7)$ is a solution to the Pell equation $X^2 - 53Y^2 = 4$. The two β values corresponding to this solution are -50 and 1 . Plugging these two values into the matrix above we get:

$$M_o = \begin{pmatrix} 1 & 7 \\ 7 & 50 \end{pmatrix} \text{ and } M_o^{-1} = \begin{pmatrix} -50 & 7 \\ 7 & -1 \end{pmatrix}.$$

The pair $(2599, 357)$ is also a solution to the above Pell equation, and the corresponding matrices are:

$$\begin{pmatrix} 50 & 357 \\ 357 & 2549 \end{pmatrix} \text{ and } \begin{pmatrix} -2549 & 357 \\ 357 & -50 \end{pmatrix}.$$

We would like to remark also that

$$M_o^2 = \begin{pmatrix} 50 & 357 \\ 357 & 2549 \end{pmatrix}.$$

In fact, M_o is one of the two fundamental automorphisms of f .

Note that the map $W \mapsto f_W$ is infinite to one because any indefinite binary quadratic form has infinite automorphism group. Any matrix in the automorphism group of f maps onto f .

Let $\mathcal{D} := \{d \in \mathbb{Z}_{>0} : d \equiv 0, 1 \pmod{4}, d \text{ is not a square}\}$. Recall the following:

Proposition 4.2 ([19]). *There is a bijection between the set of conjugacy classes of primitive hyperbolic elements in $\mathrm{PSL}_2(\mathbb{Z})$ and the set of classes of primitive binary quadratic forms of discriminant $\Delta \in \mathcal{D}$; where a hyperbolic element is called primitive if it is not a power of another hyperbolic element.*

4.1 Reduction Theory of Binary Quadratic Forms

We say that an indefinite binary quadratic form f is *reduced* if the geodesic in \mathcal{H} connecting the two real fixed points of W_f , called the *axis* of W_f and denoted by \mathfrak{a}_{W_f} , intersects with the fundamental domain

$$\mathfrak{f} := \{z \in \mathbb{H} : |z| \geq 1, |\mathrm{Re}(z)| < \frac{1}{2}\},$$

of the modular group. Remark that this definition is equivalent to the one given by Gauss in [10]. The caveat of indefinite binary quadratic forms is that the $\mathrm{PSL}_2(\mathbb{Z})$ class of such a form contains more than one reduced form as opposed to definite binary quadratic forms in which case the reduced representative is unique, see [4, Section 6.8] or [5, Section 5.6] for further discussion.

Given a primitive indefinite binary quadratic form, call $f = (A, B, C)$, of discriminant $\Delta \in \mathcal{D}$, f is reduced if $|\sqrt{\Delta} - 2|A|| < B < \sqrt{\Delta}$, the equivalence of the two definitions we use is folklore. Our aim is now to reveal the reduction method due to Gauss in terms of çarks. For this let us first note that while proving Theorem ?? we have seen that every edge of a çark may be labeled with a unique coset of the corresponding subgroup. Thus binary quadratic forms may be used to label the edges of the çark by Corollary ??.

Given a hyperbolic element W as a word in R, R^2 and S we define the length of W , $\ell(W)$, to be the total number of appearances of R, R^2 and S . For instance for $W = RSR^2S(RS)^2$, $\ell(W) = 8$. Before stating the next lemma let us recall that the classical reduction process is simply the process of acting on a non-reduced form $f = (A, B, C)$ by the matrix

$$\rho(f) = \begin{pmatrix} 0 & 1 \\ 1 & t(f) \end{pmatrix} = S(RS)^{t(f)};$$

$$\text{where } t(f) = \begin{cases} \mathrm{sign}(c) \left\lfloor \frac{b}{2|c|} \right\rfloor & \text{if } |c| \geq \sqrt{\Delta} \\ \mathrm{sign}(c) \left\lfloor \frac{\sqrt{\Delta}+b}{2|c|} \right\rfloor & \text{if } |c| < \sqrt{\Delta} \end{cases}.$$

Lemma 4.3. *Given an indefinite binary quadratic form (reduced or non-reduced), f , let W_f be a primitive hyperbolic element corresponding to f . Then*

$$\ell(W_{\rho(f)f}) \leq \ell(W_f). \quad (4.4)$$

Let us assume from now on that our çarks are embedded into an annulus, with an orientation which we will assume to be the usual one. In addition we also introduce the following shorter notation for our çarks: in traversing the spine (in either direction) if there are n consecutive Farey branches in the direction of the same boundary component, then we denote this as a single Farey component and write n on the corresponding branch, see Figure 5. We will call such çarks *weighted*.

Let us make the following

Definition 4.1. Let \mathcal{C} be a weighted çark. Edges of the spine are called *semi-reduced*. In particular, an edge on the spine of \mathcal{C} is called *reduced* if and only if it is on the either side of a Farey component which is in the direction of the inner boundary component.




Figure 6: A çark and its short form.

Remark that as we have fixed our orientation to be the usual one, there is no ambiguity in this definition. In addition note that semi-reduced edges are in one to one correspondence between the forms $f = (A, B, C)$ in a given class for which $AC < 0$. Edges labeled by binary quadratic forms $f = (A, B, C)$ for which both $A > 0$ and $C > 0$ are on a Farey component in the direction of the inner boundary. Likewise, whenever $A < 0$ as well as C the edge is on a Farey component in the direction of outer boundary of the annulus. We are now ready to describe reduction theory of binary quadratic forms in terms of çarks. We have seen that multiplication by the matrix $\rho(f)$ is, in general, the process of moving the base edge of the çark to the spine as a result of Lemma 4.3. However, this is not enough. That is, not every edge on the spine corresponds to a reduced form. Reduced forms correspond to edges where the Farey branches switch from one boundary component to the other. More precisely, we have:




Figure 7: Çark corresponding to the class represented by the form $(7, 33, -15)$. Marked edges are reduced.

Theorem 4.4. *Reduced forms in an arbitrary indefinite binary quadratic form class $[f]$ are in one to one correspondence between the reduced edges of the çark corresponding to the given class.*

As we have remarked the action of $\text{PSL}_2(\mathbb{Z})$ on binary quadratic forms is equivalent to the change of base edge on the set of çarks. Hence the above Theorem is an immediate consequence of the following:

Lemma 4.5. *Let ζ_f denote the çark associated to an arbitrary indefinite binary quadratic form f . The reduction operator $\rho(f)$ is transitive on the set of reduced edges of ζ_f .*

Let us give some examples:

Example 4.2. Let us consider the form $f = (7, 33, -15)$. It is easy to check that f is reduced. $W_f = (R^2S)^2 (RS)^2 R^2S RS (R^2S)^7 (RS)^5 = \begin{pmatrix} -38 & -195 \\ -91 & -467 \end{pmatrix}$. The trace of the class is -505 . By Gauss' theory the class $[f]$ is an element in the quadratic number field with discriminant 1509.

Example 4.3. Let $\Delta = n^2 + 4n$ for some positive integer n . Then the identity in the class group is given by the çark in Figure 8a and the corresponding form is $(-n, n, 1)$. If $\Delta = n^2 + 4$, then the identity is represented by the form $\frac{1}{n}(-n, n^2, n) = (1, n, -1)$. The corresponding çark has two Farey branches, see Figure 8b

However, one has to admit that there are very complicated çarks representing the identity of the class group. For instance, the çark corresponding to the form $(-7, 23, 16)$ has 42 Farey branches.

4.2 Ambiguous and Reciprocal forms

Let us now discuss certain symmetries of a çark. For a given çark ζ let ζ^r be the çark which is the mirror image of ζ about any line passing through the 'center' of




Figure 8

the spine (assuming that the Farey components coming out of the spine in its shorter notation that we have introduced is evenly spaced). It is easy to see that both ideal classes represented by the two çarks \mathcal{C} and \mathcal{C}^r have the same discriminant. A straightforward computation leads to the following:

Proposition 4.6. *Given a çark \mathcal{C} the binary quadratic form class represented by \mathcal{C}^r is inverse of the class represented by \mathcal{C} .*

Example 4.4. Let us consider the form $f = (-2377, 10173, 1349)$ having discriminant 116316221. The form $g = (-4027, 8915, 2287)$ is an element in the ideal class represented by this form. The corresponding çarks are shown in Figure 9. The forms are inverses of each other, see Figure 9c.




Figure 9: Two çarks inverses of one another and their product.

Recall that Gauss has defined a binary quadratic form to be ambiguous if it is equivalent to its inverse or equivalently if the corresponding equivalence class

contains (a, ka, c) for some a, c and k . Following Gauss, we define a çark \mathcal{C} *ambiguous* if \mathcal{C} and \mathcal{C}^r are isomorphic as çarks, or equivalently correspond to the same subgroup of $\mathrm{PSL}_2(\mathbb{Z})$. So from Proposition 4.6 we deduce:

Corollary 4.7. *Ambiguous çarks correspond to ambiguous forms.*

In addition to all the examples considered in Example 4.3, which represent ambiguous classes as they are of the form (a, ka, c) , let us give one more example:

Example 4.5. Consider the form $f = (3, 18, -11)$. The form is reduced and ambiguous as one immediately checks. The corresponding çark is given in Figure 10




Figure 10: Çark corresponding to the ambiguous form $f = (3, 18, -11)$.

Let us now discuss “rotational” symmetries. An oriented çark with a base edge is called *primitive* if and only if its spine is not periodic. Let \mathfrak{c}_{prim} denote the set of primitive çarks. It is easy to see that primitive hyperbolic elements⁶ in $\mathrm{PSL}_2(\mathbb{Z})$ correspond to primitive çarks or equivalently to prime geodesics in \mathbb{H} .

Corollary 4.8. *There is a one to one correspondence between the following two sets:*

$$\mathfrak{c}_{prim} \longleftrightarrow \left\{ \begin{array}{l} \mathrm{PSL}_2(\mathbb{Z}) \text{ classes of primitive} \\ \text{indefinite binary quadratic forms} \\ \text{having discriminant } \Delta \in \mathcal{D} \end{array} \right\}$$

Finally, let \mathcal{C}^m denote the mirror of a given çark, that is the çark obtained by reflecting \mathcal{C} with respect to the spine. Once again observe that both \mathcal{C} and \mathcal{C}^m have the same discriminant. In fact, an indefinite binary quadratic form say $f = (A, B, C)$ is given which is represented by the çark \mathcal{C} then the çark \mathcal{C}^m represents the form $f' = (-A, B, -C)$ and the same holds for every element in $[f]$. We conclude that both çarks represent ideal classes that have the same order in the class group.

⁶Recall that an element $M \in \mathrm{PSL}_2(\mathbb{Z})$ is said to be *primitive* if it is not a positive power of another element of the modular group.

Let W be a hyperbolic element in $\mathrm{PSL}_2(\mathbb{Z})$. In [19], Sarnak has defined W to be reciprocal if W is conjugate to its inverse. The conjugation turns out to be done by a unique element (up to multiplication by an element in $\langle W \rangle$) of order 2, and thus reciprocal elements correspond to maximal dihedral subgroups of the modular group⁷. A form $f = (A, B, C)$ is called reciprocal if $C = -A$. It is known that reciprocal hyperbolic elements correspond to reciprocal indefinite binary quadratic forms, [19]. In a similar fashion we call a çark *reciprocal* if ζ and $(\zeta^m)^r$ are isomorphic as çarks. In fact since the two operator \cdot^m and \cdot^r commute, if ζ is a reciprocal çark then so is ζ^m .

Proposition 4.9. *Reciprocal forms correspond to reciprocal çarks.*




Figure 11: The graph $\mathcal{F}/\langle S, R^2SR \rangle$




Figure 12: The graph $\mathcal{F}/\langle RSR^2, S(RSR^2)S \rangle$

Example 4.6. Consider the form $f = (-8, 11, 8)$. The corresponding hyperbolic element in $\mathrm{PSL}_2(\mathbb{Z})$ is $\begin{pmatrix} 101 & -192 \\ -192 & 365 \end{pmatrix}$. The corresponding çark is shown in Figure 13, where it is easy to see that ζ and $(\zeta^m)^r$ are same.

Example 4.7 (Reciprocal Identities). The forms $f = (1, n^2, -1)$ already appeared in Example 4.3 are reciprocal and represent identity in the class group. Conversely, given a form, which is both reciprocal and represents identity has to be of this form. Hence we conclude that identity elements that are at the same time reciprocal appear only for discriminants $n^2 + 4$ and cenversely. Note also that all such forms come from the word $(R^2S)^n(RS)^n$. The çarks of reciprocal identities are in Figure 8b

⁷Remember that çarks correspond to maximal \mathbb{Z} subgroups of $\mathrm{PSL}_2(\mathbb{Z})$.




Figure 13: Çark corresponding to the reciprocal form $f = (-8, 11, 8)$.

4.3 Miscellany

Binary quadratic forms is a central and classical topic and have connections to diverse fields. Here we touch upon some of these.

4.3.1 Computational Problems

There are several important computational problems related to çarks, in connection with the class number problems in the indefinite case. The most basic invariant of a çark is the length of its spine. The (absolute) trace of the associated matrix is another, much subtler invariant. We don't know any practical means to compute the trace from the combinatorial input (bracelet) of a çark, other than explicitly computing the associated matrix. Similarly, given two çarks, we don't know any direct way of saying whether they are of the same trace or not. The problem of listing çarks of the same trace is equivalent to the problem of computing class numbers. Also, the Gauss product on classes of forms defines an abelian group structure on the set çarks of the same trace, namely the *class group*. We have not been successful in our attempts to reach to a new understanding of class groups in terms of the graphical representation of their elements by çarks.

4.3.2 Closed geodesics on the modular surface.

Let us note in passing that primitive çarks parametrize closed geodesics on the modular curve, and so çarks are closely connected to symbolic dynamics on the modular curve. [13], encoding of geodesics, and Selberg's trace formula. [21]

4.3.3 The Markoff number of an indefinite binary quadratic form.

There is an arithmetic invariant of indefinite binary quadratic forms called the Markoff value $\mu(F)$ which is defined as

$$\mu(F) := \frac{\sqrt{\Delta(F)}}{m(f)}, \text{ where } m(f) := \min_{(x,y) \in \mathbb{Z}^2 \setminus \{(0,0)\}} |F(x,y)|$$

Alternatively one can run over the class of F and compute the minima of equivalent forms at a fixed point p_0 , for example $(x, y) = (0, 1)$. Hence the choice of this fixed point p_0 defines a function on the set of edges of the associated çark, and the Markoff value of the form is the maximal value attained by this function defined on the çark. There are also çarks associated to Markoff irrationalities which we call *Markoff çarks*.




Figure 14: Minimum edges of $\mathcal{F}/\text{Aut}^+\{(1, 0, -2)\}$.

Referanslar enddocumnet in altinda kaldi!!!!!!

Acknowledgements. This research has been funded by the TÜBİTAK grant 110T690. The first named author has also been funded by a GSU Research Fund Grant during the preparation of the current paper.

References

- [1] *The on-line encyclopedia of integer sequences*, published electronically at <http://oeis.org/>.
- [2] G. Belyi, *On galois extensions of a maximal cyclotomic field*, Math. USSR Izv. **14**, No.2 (1980), 247–256.
- [3] Salwa Bouallègue and Mongi Naimi, *On primitive words*, Int. J. Algebra **4** (2010), no. 13-16, 693–707. MR 2725470 (2011m:68220)
- [4] Johannes Buchmann and Ulrich Vollmer, *Binary quadratic forms*, Algorithms and Computation in Mathematics, vol. 20, Springer, Berlin, 2007, An algorithmic approach. MR 2300780 (2008b:11046)

- [5] Henri Cohen, *A course in computational algebraic number theory*, Graduate Texts in Mathematics, vol. 138, Springer-Verlag, Berlin, 1993. MR 1228206 (94i:11105)
- [6] John H. Conway, *The sensual (quadratic) form*, Carus Mathematical Monographs, vol. 26, Mathematical Association of America, Washington, DC, 1997, With the assistance of Francis Y. C. Fung. MR 1478672 (98k:11035)
- [7] Merve Durmuş, *Farey Graph and Binary Quadratic Forms*, Master's thesis, İstanbul Technical University, Turkey, 2012.
- [8] J.-P. Duval, *Generation d'une section des classes de conjugaison et arbre des mots de Lyndon de longueur bornée*, Theoret. Comput. Sci. 60 (1988) 255-283.
- [9] H. Fredricksen and I. J. Kessler, *Lexicographic compositions and de Bruijn sequences*, Journal of Combinatorial Theory 22 (1977), 17-30.
- [10] Carl Friedrich Gauss, *Disquisitiones arithmeticae*, Translated into English by Arthur A. Clarke, S. J, Yale University Press, New Haven, Conn., 1966.
- [11] Ernesto Girondo Gabino Gonzalez-Diez Introduction to Compact Riemann Surfaces and Dessins d'Enfants London Mathematical Society Student Texts 79
- [12] Svetlana Katok, *Reduction theory of Fuchsian groups.*, Math. Ann. **273** (1986), 461-470 (English).
- [13] Svetlana Katok and Ilie Ugarcovici, *Symbolic dynamics for the modular surface and beyond*, Bull. Amer. Math. Soc. (N.S.) **44** (2007), no. 1, 87-132.
- [14] F. Klein, *Über die transformation elfter ordnung der elliptischen functionen*, Mathematische Annalen **15** (1879), no. 3-4, 533-555.
- [15] Ravi S. Kulkarni, *An arithmetic-geometric method in the study of the subgroups of the modular group*, Amer. J. Math. **113** (1991), no. 6, 1053-1133.
- [16] S. Lando and A. Zvonkin, *Graphs on surfaces and their applications*, Encyclopaedia of Mathematical Sciences, Springer, 2004.
- [17] P. A. MacMahon, *Applications of a Theory of Permutations in Circular Progression to the Theory of Numbers*, Proc. London Math. Soc. **S1-23**, no. 1, 305.
- [18] Jin Ho Kwak, Roman Nedela, Graphs and their Coverings, <http://www.savbb.sk/nedela/graphcov.pdf>
- [19] P. Sarnak, *Reciprocal geodesics*, Analytic number theory, Clay Math. Proc., vol. 7, Amer. Math. Soc., Providence, RI, 2007, pp. 217-237.
- [20] Joe Sawada *Generating Bracelets in Constant Amortized Time* SIAM Journal on Computing archive 31 Issue 1, 2001, 259 - 268.

- [21] D. Zagier, *New points of view on the selberg zeta function*, Proceedings of Japanese-German Seminar, 2002.
- [22] D. Zagier, *Zetafunktionen und quadratische Körper: eine Einführung in die höhere Zahlentheorie*, Hochschultext, Springer, Berlin, Heidelberg, New York, 1982